



Published in final edited form as:

*Clin Geriatr Med.* 2014 August ; 30(3): 421–442. doi:10.1016/j.cger.2014.04.001.

## Dementia and Cognitive Impairment: Epidemiology, Diagnosis, and Treatment

Julie Hugo, MD and Mary Ganguli, MD, MPH

### Synopsis

Clinicians can diagnose the syndromes of dementia (major neurocognitive disorder) and mild cognitive impairment (mild neurocognitive disorder) based on history, examination, and appropriate objective assessments, using standard criteria such as DSM-5. They can then diagnose the etiological subtypes of these syndromes using standard criteria for each of them. Brain imaging and biomarkers are gaining ground for the differential diagnoses among the different disorders. Treatments for the most part are still symptomatic.

### Keywords

Neurocognitive disorder; Mild Cognitive Impairment (MCI); Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5); National Institute on Aging- Alzheimer's Association (NIA-AA) Guidelines; diagnosis; risk factors; biomarkers; Alzheimer's disease

## INTRODUCTION

When older patients and their families report symptoms of “memory loss,” experienced clinicians know that these concerns refer to a range of cognitive abilities or to general cognitive decline, and not just memory. However, some degree of cognitive slowing is typical of normal aging.

The clinician's first challenge, therefore, is to identify the cognitive changes that are clinically significant. **Dementia** is typically diagnosed when acquired cognitive impairment has become severe enough to compromise social and/or occupational functioning. **Mild cognitive impairment (MCI)** is a state intermediate between normal cognition and dementia, with essentially preserved functional abilities.

We will describe these entities and their diagnoses using the framework of the recently published fifth edition of the American Psychiatric Association's Diagnostic and Statistical Manual (DSM-5) (Table 1).<sup>1</sup> Briefly, the DSM-5 diagnosis of Major Neurocognitive

---

© 2014 Elsevier Inc. All rights reserved.

Corresponding Author: Mary Ganguli, MD, MPH, **Professor of Psychiatry, Neurology, and Epidemiology**, University of Pittsburgh, Pittsburgh, PA, WPIC, 3811 O'Hara Street, Pittsburgh, PA 15213, GanguliM@upmc.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disorder, which corresponds to dementia, requires substantial impairment to be present in one or (usually) more cognitive domains. The impairment must be sufficient to interfere with independence in everyday activities. The diagnosis of Mild Neurocognitive Disorder, corresponding to MCI, is made when there is modest impairment in one or more cognitive domains. The individual is still independent in everyday activities, albeit with greater effort. The impairment must represent a decline from a previously higher level and should be documented both by history and by objective assessment. Further, the cognitive deficits must not occur exclusively in the context of a delirium or be better explained by another mental disorder.

The clinician's second challenge is to determine the cause(s) of the cognitive impairment, i.e., to identify the underlying etiology. DSM-5 also provides diagnostic criteria for the most common etiologic subtypes of the neurocognitive disorders in all age groups. Here we focus on the neurocognitive disorders of older adults.

### **Impact of Dementia**

Neurocognitive disorders, particularly major neurocognitive disorders (dementias), have tremendous consequences for individuals, their families, the healthcare system, and the economy. In the United States, Alzheimer's disease (AD) is a leading cause of death,<sup>2</sup> hospital admissions, skilled nursing facility admissions, and home health care. The costs of health services and the informal costs of unpaid caregiving for individuals with dementia are high and growing. Family caregivers also experience increased emotional stress, depression, and health problems.<sup>3</sup> In absolute numbers, 35.6 million people worldwide were estimated to be living with dementia in 2010, a number expected to reach 115.4 million people by 2050.<sup>4</sup>

### **Dementia in the Population**

Prevalence, defined as the proportion of people with an illness in a given population at a given time, is an index of the burden of disease in the population. Incidence is the rate at which new disease occurs in a given population, i.e., the proportion of new cases in that population over a given period of time. Incidence is therefore an index of the risk of disease in that population. Prevalence is a function of both incidence and duration. Since most dementias are not curable, their duration reflects how long individuals live with their dementia. Thus, the public health burden of dementia depends both on the development of new cases and on the survival of those cases after onset; holding incidence constant, groups with longer life expectancy will have higher prevalence.

### **Prevalence**

Prevalence of dementia increases exponentially with increasing age<sup>5</sup>, and doubles every five years of age after age 65. In higher income countries, prevalence is 5–10% in those aged 65+ years, usually greater among women than among men, in large part because women live longer than men. Within the US, higher prevalence has been reported in African American and Latino/Hispanic populations than in White nonHispanic populations. Global systematic reviews and meta-analyses suggest that prevalence of dementia is lower in sub-Saharan Africa and higher in Latin America than in the rest of the world [Table 2]. The prevalence of

MCI is at present difficult to pin down as it depends on the precise definitions and subtypes of MCI being studied.<sup>6</sup>

Life expectancy is increasing across the planet, with population aging rising the most rapidly in the low- and middle-income countries, where the prevalence of dementia is therefore expected to increase.<sup>7</sup> Emerging studies suggest that prevalence may be leveling off or even decreasing in the high-income countries.<sup>8,9</sup>

### Incidence

The incidence of dementia increases steadily until age 85 or 90, and then continues to rise but less rapidly. It is either similar in men and women or slightly higher in women. Annual age-specific rates ranged from 0.1% at age 60–64 to 8.6% at age 95.<sup>5,18</sup>

### Risk and Protective Factors

Risk factors are factors associated with an elevated incidence rate of disease, higher odds of developing disease, or earlier onset of disease, depending on the type of statistical analysis that is performed. Protective factors represent the converse. An observed risk factor does not necessarily cause disease; a protective factor does not necessarily prevent disease and almost certainly will not treat the disease. The observed effects can potentially reflect selection or survival bias or confounding, or sometimes reverse causality. They may also depend on the timing and duration of exposure to the factor, with mid-life often being the critical period.

### Demographic risk factors

Increasing age is not only the strongest risk factor for dementia, but also the only risk factor consistently identified after the eighth decade of life. While prevalence is consistently higher among women, incidence is not; thus, the higher prevalence may largely be a function of longer life-expectancy in women. Lower educational levels have been found associated with higher prevalence. Within the US, prevalence has been reported as elevated in African American and Latino populations; some authors have attributed these findings to lower education and higher cardiovascular morbidity in those populations.

### Genetic factors

Few dementias are caused by deterministic autosomal dominant genes; these are discussed later under the specific disorders. While several genes have been identified as increasing susceptibility for Alzheimer's disease, the best-established is the Apolipoprotein E (*APOE*) polymorphism on Chromosome 19. The *APOE*\*4 allele, associated with higher risks of hypercholesterolemia and heart disease, is also associated with dementia due to Alzheimer's and Parkinson's diseases, Dementia with Lewy Bodies, vascular dementia, and frontotemporal dementia in men.<sup>19,20,21,22,23</sup> Individuals homozygous for *APOE*\*4 are at greater risk of dementia than those who are heterozygous. The *APOE*\*2 appears to have a protective effect. *APOE*\*4 is a risk factor, not a diagnostic marker for Alzheimer's disease. It is neither necessary nor sufficient for diagnosis, and its effect on risk appears to wear off by the eighth decade, i.e. individuals who are older than 80 years, *APOE*\*4 positive, and do not yet have dementia, are at no greater risk of developing dementia than those who are *APOE*\*4 negative.

## Medical risk factors

Cardiovascular disease is increasingly recognized as not just a risk factor for vascular dementia but also for degenerative dementias, particularly AD. Heart disease has been associated with both dementia of the Alzheimer's type, and vascular dementia.<sup>24</sup> Risk factors in midlife, including hypertension, high cholesterol, high body mass index (BMI), and diabetes mellitus are associated with increased risk of dementia in late life, demonstrating the importance of risk exposures decades earlier.<sup>2526</sup> Heart failure and atrial fibrillation are risk factors for cognitive impairment and dementia.<sup>272829</sup> Cardiac disease can cause or worsen cerebral hypoperfusion, creating a cellular energy crisis setting off a cascade of events leading to the production of toxic proteins.<sup>30</sup> In cognitively normal older adults, elevated pulse pressure has recently been found associated with alterations in biomarkers suggestive of AD.<sup>31</sup>

Inflammation and alterations in inflammatory markers [interleukins, cytokines, C-reactive protein] have been reported in Alzheimer's and vascular dementias.<sup>3233</sup> Multiple mechanisms have been proposed for the role played by inflammation in the neuropathology of AD.<sup>343536</sup>

Obstructive sleep apnea, associated with hypertension, heart disease, stroke risk<sup>37</sup> and white matter change,<sup>38</sup> is also associated with an increased risk of dementia.<sup>39</sup>

Stroke increases risk of dementia.<sup>4041</sup>

## Psychiatric risk factors

Depression has a complex and likely bi-directional association with dementia. Recurrent major depression in earlier adulthood appears to increase risk of dementia in later life.<sup>42</sup> Depression with late life onset is believed to be an early sign of the vascular or degenerative disease causing the dementia.<sup>4344</sup> Late-life anxiety is associated with cognitive impairment and decline.<sup>45</sup> Post-traumatic stress disorder has been reported as increasing risk of dementia.<sup>46</sup> Lifelong traits of harm avoidance and lesser sense of purpose have been reported as harbingers of AD.<sup>47</sup>

## Head injury

Head injury is associated with increased risk of dementia, in particular AD,<sup>48</sup> and the severity of injury appears to heighten this risk.<sup>4950</sup> Neurocognitive disorders can occur immediately after a traumatic brain injury or after the recovery of consciousness at any age.<sup>1</sup> However, chronic traumatic encephalopathy (previously termed dementia pugilistica) is diagnosed years after repeated concussive or subconcussive blows to the head, with a clinical presentation similar to AD or frontotemporal lobar degeneration.<sup>51</sup>

## Lifestyle and environmental risk factors

Many environmental and occupational exposures have shown varying associations with neurodegenerative diseases.<sup>52</sup> Smoking has been associated with an elevated risk of dementia;<sup>53</sup> although some studies have found an apparent protective effect which could reflect survival bias (competing risks)<sup>54</sup> or possibly cholinergic action as also seen in

Parkinson's disease.<sup>55</sup> Heavy consumption of alcohol increases odds of developing dementia.<sup>56,57</sup> Parkinson's disease risk is associated with exposure to pesticides, for which a molecular mechanism has been established.<sup>58</sup>

### Protective Factors

Protective factors are those associated with a reduced incidence rate or reduced odds of dementia, or with delayed onset of dementia.

The concept of "reserve" was proposed to explain why some individuals remain cognitively intact despite the presence of neuropathology typically associated with dementia.<sup>59</sup> Brain reserve refers to structural capacity and integrity of the brain (e.g. brain mass, preserved large neurons),<sup>60</sup> whereas cognitive reserve refers to its functional capacity, specifically the ability to utilize alternative neural networks and compensatory strategies.<sup>61</sup>

### Education and cognitive activity

Where educational opportunities are universal, higher education may reflect innate reserve; the process of education may also promote the development of reserve through mechanisms such as increased dendritic branching. Education may also reflect general socioeconomic status and thus also represent quality of environmental factors like nutrition, or health care. Regardless of mechanism, higher education is associated with lower prevalence of dementia.<sup>62</sup>

Bilingualism has been associated with delayed onset of dementia,<sup>63</sup> independent of education, and may specifically protect against declines in attention and executive functioning.<sup>64</sup>

Cognitive activity: Lifelong occupations that do not require higher education or skilled vocational training appear to be associated with a higher risk of dementia.<sup>65,66</sup> Several popular leisure activities have been found associated with lower risk of dementia.<sup>67</sup> Cognitively stimulating activities appear to have both protective and enhancing effects on cognition.<sup>68</sup>

### Pharmacological factors

Several therapies for other conditions have been found in long-term observational studies to be associated with a reduced risk of dementia. However, in clinical trials, these drugs have not been found to prevent dementia. Timing and duration of the exposure might partly explain these discrepancies, as the protective effects were seen with prolonged use multiple years before dementia onset. While some studies have found a protective relationship with the use of non-steroidal anti-inflammatory drugs (**NSAIDs**),<sup>69,70,71</sup> a 2005 meta-analysis determined that many of the positive results seen in the 25 studies reviewed were due to various forms of bias.<sup>72</sup> Despite mixed reports of the effects of the lipid-lowering HMG Co-A reductase inhibitors ("**statins**"), pooled results in a recent review as well as in a 2013 meta-analysis indicate a protective effect against dementia.<sup>73,74</sup>

As regards **estrogen** therapy, the WHIMS trial in older women showed no protection and possibly an increased risk from combination hormone therapy.<sup>75,76</sup> A 2013 meta-analysis of

any use versus no use concluded that hormones had no effect on dementia.<sup>77</sup> However, a long-term observational study suggests that the timing of hormone therapy, at menopause, may be the critical factor.<sup>78,79,80</sup>

### Lifestyle factors

As with cardiovascular disease, population based studies have found that mild to moderate alcohol consumption is associated with reduced risk of cognitive impairment and dementia.<sup>56,57,81,82</sup> Adherence to a Mediterranean diet was associated with better cognitive functioning, lower rates of cognitive decline and a reduced risk of AD.<sup>83</sup> High physical activity levels are associated with reduced risk of neurodegenerative diseases.<sup>84</sup> Smoking shows a consistently protective effect against Parkinson's disease, potentially involving nicotine effects on cholinergic receptors.<sup>55</sup>

Elderly women with larger social networks,<sup>85</sup> and who participate in mental, social or productive activities, have shown lower incidence of dementia.<sup>86</sup> Social, mental and physical lifestyle components appear important,<sup>87</sup> although reverse causality cannot be ruled out, given that the neuropathology often begins decades before symptom onset.

## CLINICAL ASSESSMENT

Assessment is easier for the clinician when patients themselves and/or their family members or other caregivers spontaneously express concern about cognitive difficulties. In other cases, neither patient nor caregiver report concerns but may acknowledge them if asked. Alternatively, the patient denies any difficulty, and there is no reliable informant, but the clinician observes cognitive impairment, this being more likely when the clinician knows the patient well. Sometimes the clinician, or a member of the clinic staff, is tipped off by a patient forgetting to keep appointments or fill prescriptions, or being confused by simple instructions. Experienced clinicians also recognize vagueness and evasiveness in patient's responses, or catastrophizing of minor problems, as clues to failing cognition. Occasionally, the initial complaint will not be of cognitive loss but of changes in mood or behavior, such as apathy, anxiety, or depression.

### Subjective assessment

Whenever possible, history should be obtained both from the patient and also from a family member, caregiver, or other reliable informant. In some mild cases, the patient is more aware of the deficits than the relative. The clinician should focus on changes in cognitive functioning as manifested in everyday activities. Early deficits are frequently noted in managing finances and medications. Over-learned, routine activities may be preserved but problems may be occurring in problem-solving, multi-tasking, and dealing with new situations. Table 3 lists the cognitive domains recognized by DSM-5 and everyday activities that tap those domains.<sup>1</sup>

Many patients and families accept cognitive decline as part of normal aging, and will declare themselves normal on the grounds that they are no worse than others their age. Others may magnify the significance of relatively minor changes and express fears of developing AD. Despite the value of systematic and disciplined clinical observations, clinicians' impressions

can be clouded by personal expectations of what is normal for a given patient. Subjective concerns alone are therefore insufficient for diagnosis.

### Objective assessment

The objective assessment requires the administration of one or more standardized tests. Neuropsychological assessment of specific cognitive domains is preferred both for detecting mild impairments and for differential diagnosis (Table 4). Details of such assessments are beyond the scope of this chapter but are readily available from other sources. If neuropsychological assessment is unavailable, objective testing can consist of a global screening scale, such as the well-known Mini-Mental State Examination (MMSE),<sup>88</sup> the Montreal Cognitive Assessment (MoCA),<sup>89</sup> or the Mini-Cog.<sup>90</sup> Such tests are usually sensitive enough to detect dementia but not necessarily MCI. It is critically important that a patient's test performance be interpreted in accordance with norms for that patient's age and educational level, and preferably for his/her cultural/linguistic group and region as well.

### Additional assessments

A general physical and neurological examination, and appropriate laboratory investigations, should be performed both to rule out treatable causes of cognitive impairment (even if they only partially explain the impairment) and to aid in differential diagnosis.

We present here the etiologic subtypes most likely to be seen in geriatric psychiatry settings.

## Alzheimer's Disease

Alzheimer's disease (AD) is the single most common neurodegenerative disease, characterized by progressive loss of synapses and neurons, with the accumulation of amyloid plaques, and neurofibrillary tangles, and prominent cholinergic deficits. It is typically diagnosed in the eighth or ninth decades of life, but early onset forms of the disease may be diagnosed as early as the fifth decade. Average duration of survival is about 10 years after the onset of dementia, but varies widely depending on the age of onset, the severity of cognitive impairment, the presence of comorbid diseases, and other factors.<sup>91,92</sup>

In DSM-5,<sup>1</sup> AD is listed an etiologic subtype of both Major and Mild Neurocognitive disorders. Criteria for this subtype harmonize with the latest expert guidelines for Dementia and MCI due to AD, as published by the National Institute on Aging – Alzheimer Association (NIA-AA) Work Group.<sup>93</sup> Unlike the NIA-AA guidelines, the DSM-5 criteria are intended primarily for clinical rather than research use, and do not include pre-clinical AD. The diagnosis of Dementia (Major NCD) in Alzheimer's disease requires evidence of decline to the level of substantial impairment in at least two cognitive domains, one of which must be memory. To diagnose MCI (Mild NCD) in Alzheimer's disease, decline to the level of modest impairment must be observed in memory and potentially in additional domains as well. The cognitive decline should be of insidious onset with gradual and steady progression. Impairments in memory and executive functions typically develop earlier in the disease course, while impairments in visuoconstructional/ perceptual- motor functions, language functions, and social cognition occur later. However, non-amnestic presentations also do occur. Depression and apathy may occur throughout the clinical spectrum. In the

middle to later stages, psychotic features, irritability, agitation, combativeness and wandering may occur, and very late in the illness, gait disturbances, dysphagia, incontinence, myoclonus and seizures may be evident.<sup>1</sup>

For neurocognitive disorders to be attributed to Probable AD, there should either be evidence of autosomal dominant familial AD, or no evidence of mixed etiology, i.e., no other contributing neurological, psychiatric, or systemic disorder that could explain the cognitive decline. Otherwise, the diagnosis of Possible AD is appropriate.<sup>1</sup>

### Genetics of AD

Autosomal dominant mutations which cause rare cases of early onset familial AD are the amyloid precursor protein (APP) gene on Chromosome 21, the Presenilin 1 (PS1) gene on Chromosome 14, and the Presenilin 2 (PS2) gene on Chromosome 1. Individuals with Down's Syndrome, caused by Trisomy 21, inevitably develop Alzheimer's neuropathology if they live long enough. The *APOE\*4* gene, as noted under Risk Factors, increases risk of dementia but is not diagnostic.

### Biomarkers for AD

Evidence of cerebral amyloid deposition, such as positron emission tomographic (PET) brain scans with amyloid tracers, and reduced levels of amyloid beta 42 in the cerebrospinal fluid (CSF), have been proposed as research biomarkers. Evidence of neuronal injury, such as hippocampal atrophy on magnetic resonance imaging (MRI) brain scan, temporoparietal hypometabolism on fluorodeoxyglucose (FDG) PET scans, and elevated total tau and phospho-tau levels in CSF, are less specific to AD but have also been proposed as research biomarkers.<sup>1</sup> They have not been officially validated or approved for clinical diagnostic use.

Unlike DSM-5, the NIA-AA guidelines also describe a stage of asymptomatic preclinical Alzheimer's disease, in which the pathology of the disease is present as evidenced by biomarkers and subclinical cognitive decline detectable only by objective testing.<sup>94</sup>

### Vascular Dementia (Vascular Neurocognitive Disorder)

In Major and Mild Vascular Neurocognitive Disorders,<sup>1</sup> the cognitive deficits are principally attributed to cerebrovascular disease. Various referred to as arteriosclerotic dementia, multi-infarct dementia, vascular cognitive impairment and vascular cognitive disorder,<sup>95</sup> it is the second most common cause of dementia and is frequently present in combination with AD ("mixed dementia"). It can result from both large and small vessel disease, with the location of the lesions more important than the volume of destruction.<sup>95</sup> Given the variability of lesions and locations, the presenting symptoms and time course are often variable. The progression of the neurocognitive decline can be in an acute stepwise pattern, show a more gradual pattern, or can be fluctuating or rapid in its course.<sup>1</sup>

To diagnose vascular neurocognitive disorder, there should either be a clear history of stroke or transient ischemic attacks temporally related to the cognitive decline, or neurological deficits consistent with sequelae of previous strokes. Cognitive decline is usually seen in the domains of complex attention and executive functions. Gait disturbance, urinary symptoms,

and personality or mood changes (including emotional lability) are common.<sup>1</sup> The depression associated with vascular neurocognitive disorder may have a late-life presentation and be coupled with psychomotor slowing and executive dysfunction, the so called vascular depression.<sup>96</sup>

### Neuroimaging

Neuroimaging (CT or MRI) evidence of significant parenchymal injury attributable to cerebrovascular disease can include one or more large vessel infarcts, a single large or strategically located infarct or hemorrhage, extensive lacunar infarcts outside the brainstem, or extensive white matter lesions.

### Genetics

There are rare autosomal dominant cerebrovascular disorders, such as CADASIL (cerebral autosomal-dominant arteriopathy with subcortical infarcts and leukoencephalopathy), which is a form of hereditary stroke caused by Notch-3 mutations on Chromosome 19.

## Frontotemporal lobar degeneration (Fronto-temporal Dementia)

Frontotemporal dementia (FTD), the third most prevalent degenerative dementia, is characterized by prominent atrophy of the frontal and temporal lobes, with the predominant neuropathological proteins containing inclusions of hyperphosphorylated tau or ubiquitin protein.

With mean onset in the sixth decade, FTD is a common cause of early-onset dementia although 20–25% of individuals with this disorder are over age 65.<sup>1</sup> The duration of survival is 6–11 years after symptom onset, and 3–4 years after diagnosis.<sup>97</sup> With insidious onset and gradual progression, the clinical subtypes (behavioral and language variants) of FTD correspond to specific areas of brain atrophy.

In the **Behavioral Variant**, changes in personality and behavior are most prominent, with loss of interest in personal affairs and responsibilities, social withdrawal, loss of awareness of personal hygiene, and socially disinhibited behavior.<sup>97</sup> Perseverative or compulsive motor behaviors, as well as hyperorality and dietary changes may also be evident.<sup>1</sup> These patients are often initially seen in psychiatric settings and can be misdiagnosed as major depressive or bipolar disorder. In addition to the Behavioral Variant, there are three **Language Variants**: (i) the **Semantic** type which appears as a fluent aphasia with impoverished content and paraphasic errors, with intact syntax and prosody; emotional blunting, loss of empathy, and rigid behaviors may also be seen,<sup>97</sup> (ii) **Progressive Nonfluent Aphasia**, and (iii) the **Logopenic** subtype.

### Genetics

In familial FTD, mutations have been associated with genes encoding proteins affecting a number of fundamental cellular functions, including microtubule-associated protein tau (MAPT), granulin (GRN), C9ORF72, transactive response DNA-binding protein of 43 kDa, valosin-containing protein, chromatin modifying protein 2B, and fused in sarcoma protein.

## Neuroimaging

Structural MRI or CT can show distinct patterns of regional cortical atrophy which correlate with the clinical variants of FTD.

## Dementia with Lewy Bodies

Dementia with Lewy bodies (DLB) is the second most common neurodegenerative dementia. The underlying disease is primarily characterized by alpha-synuclein misfolding and aggregation within the pathognomonic Lewy bodies, which are also found in Parkinson's disease. Onset of symptoms is between the sixth and ninth decades, and average survival is 5–7 years.<sup>1</sup>

With insidious onset and gradual progression, the cognitive deficits are most prominent in the domains of attention, visuospatial and executive functioning. Additional core features include fluctuating cognition, recurrent visual hallucinations, and parkinsonism.<sup>1</sup> The key distinction between DLB and dementia of Parkinson's disease is based on the temporal sequence of the cognitive impairment and the movement disorder. In DLB, cognitive impairment precedes the onset of parkinsonism, while in the latter, the cognitive impairment occurs in the context of established Parkinson's disease.

Suggestive features of DLB include REM sleep behavior disorder and severe neuroleptic sensitivity. Low dopamine transporter (DaT) uptake in basal ganglia demonstrated by SPECT or PET imaging has been proposed as a suggestive feature. Supportive clinical features include repeated falls and syncope, transient and unexplained loss of consciousness, severe autonomic dysfunction, hallucinations in other modalities, systematized delusions and depression.<sup>98</sup>

## Neuroimaging

To help differentiate Lewy-body related dementias (DLB and Dementia in Parkinson's disease) from other dementias, dopamine transporter (DaT) PET scans may be useful. Generalized low uptake on SPECT and fluorodeoxyglucose PET with reduced occipital activity also suggests DLB. Additional testing supportive of DLB include low uptake MIBG myocardial scintigraphy, suggesting synaptic denervation, as well as prominent slow wave activity on EEG with temporal lobe transient sharp waves.<sup>98</sup>

## Neurocognitive Disorders due to Parkinson's Disease

These disorders are diagnosed when there is gradual cognitive decline in the presence of a well-established diagnosis of Parkinson's disease. Over the course of their disease, approximately 75% of individuals with Parkinson's disease will develop a major neurocognitive disorder.<sup>1</sup> The pattern of cognitive deficits is variable but often affects the executive, memory, and visuospatial domains, with a slowing of information processing that suggests a "subcortical" picture. Associated features include psychiatric symptoms such as depressed or anxious mood, apathy, hallucinations, delusions, or personality change, as well as rapid eye movement sleep behavior disorder and excessive daytime sleepiness.<sup>1</sup>

## Neurocognitive Disorder due to Huntington's Disease

Huntington's disease is a neurodegenerative disease caused by an autosomal dominant mutation consisting of CAG repeats on Chromosome 4. The neurotoxic Huntingtin (HTT) protein begins by damaging the striatum of the basal ganglia but eventually affects the entire brain. Although adult onset Huntington's disease usually manifests in the fourth or fifth decades, patients have a median survival of 15–20 years after diagnosis, and can thus present to geriatric services. A few patients develop their first symptoms at older ages in the absence of a family history. Progressive cognitive impairment to eventual dementia is inevitable. Although cognitive deficits (executive function) and behavioral symptoms (depression, anxiety, apathy, obsessive-compulsive symptoms, and psychosis) often emerge before the motor abnormalities (bradykinesia and chorea), clinical diagnosis is rarely made on the basis of cognitive symptoms alone. A family history of the disease should alert clinicians to the possibility, and genetic testing for the HTT mutation is diagnostic.<sup>1</sup>

## Neurocognitive Disorder due to Prion Disease

These are neurocognitive disorders due to spongiform encephalopathies caused by transmissible misfolded protein particles called prions. The human prion disorders include kuru, sporadic Creutzfeldt-Jacob disease (CJD), familial CJD, iatrogenic CJD, Gerstmann-Sträussler-Scheinker disease, fatal insomnia, and new variant CJD. Human transmission has been reported due to infected growth hormone injection and corneal transplantation; cross-species transmission is exemplified by bovine spongiform encephalopathy ("mad cow disease.") These illnesses progress rapidly and combine neurocognitive decline and motor features such as myoclonus and ataxia. Variant CJD may present with low mood, withdrawal, and anxiety. Individuals are typically diagnosed in their seventh and eighth decades, and the course is rapidly progressive, with survival typically under one year.<sup>99</sup> Diagnosis can only be confirmed by biopsy or autopsy. However, MRI scanning with diffusion weighted imaging or fluid-attenuated inversion recovery may show multifocal gray matter hyperintensities in the subcortical and cortical areas. Tau or 14-3-3 protein may be found in the cerebrospinal fluid; characteristic triphasic waves may be seen on the electroencephalogram. Genetic testing may be useful in the 15% of cases who have a family history suggesting an autosomal dominant mutation.<sup>1</sup>

## TREATMENT

### Etiology-specific treatment

If a neurocognitive disorder is diagnosed as wholly or partly due to a treatable condition, treatment specific to that condition is clearly the first line of defense. At this time, no disease-modifying therapies are available for any of the neurodegenerative diseases. However, symptomatic and supportive treatments are usually of value.

### Symptomatic Treatment

**Cholinesterase inhibitors** increase cholinergic transmission at the synaptic cleft, potentially benefiting patients with cholinergic deficits as in AD. Three such drugs are currently available in the US: donepezil, rivastigmine, and galantamine. For dementia due to AD, a

systematic review determined that all three drugs are comparable in efficacy and, on average, provide modest improvements in cognitive function and everyday activities and behavior in Alzheimer's disease.<sup>100</sup> Although approved for use in severe dementia, many practitioners question its value in advanced disease. Evidence is mixed on the effects of these drugs on long-term outcomes, e.g., slowing of the rate of decline in everyday functions, and delay of institutionalization.<sup>101</sup>

Rivastigmine is also approved for dementia in Parkinson's disease. A large double-blind placebo-controlled trial of rivastigmine showed meaningful improvements in cognition and everyday functioning.<sup>102103</sup>

While there is expert consensus that cholinesterase inhibitors are more effective in DLB than in AD, for both cognitive and behavioral effects,<sup>104</sup> evidence from large controlled trials is lacking.

In vascular dementia, evidence is mixed for the cholinesterase inhibitors. They are often prescribed in vascular dementia because of the frequent co-occurrence of cerebrovascular and neurodegenerative disease.<sup>105</sup>

In frontotemporal dementia, there is no convincing evidence of benefits from these drugs, and there are reports that they worsen behavior symptoms.<sup>106107</sup>

There is inadequate evidence on the use of cholinesterase inhibitors in other neurocognitive disorders.

In contrast, a systematic review has found minimal evidence of benefit from these drugs in mild cognitive impairment, either with symptom relief or delay in progression to dementia. Further, this weak evidence was overwhelmed by the risk of adverse effects, particularly gastrointestinal effects.<sup>108</sup>

### **NMDA receptor antagonist**

One such agent, memantine, is approved for the treatment of moderate to severe dementia due to Alzheimer's disease. It is believed to be neuroprotective against excitotoxicity in the cortex and hippocampus. An advantage of memantine is that it is well tolerated.

A systematic review showed that memantine had a small beneficial effect on cognition at six months in moderate to severe AD, marginal effect on mild to moderate AD, and a small but clinically undetectable effect in mild to moderate vascular dementia.<sup>109</sup>

In frontotemporal dementia, memantine has shown mixed results.<sup>106</sup> There is preliminary evidence of benefits in DLB and Dementia in Parkinson's disease; however, there have been reports of worsening delusions and hallucinations in DLB.

### **Serotonergic agents**

SSRI antidepressants can produce benefits for behavioral /psychiatric symptoms in frontotemporal dementia, without concomitant improvements in cognition.<sup>110</sup>

## Dopamine blocking agents

Neuroleptic (antipsychotic) drugs should be prescribed in dementia with due attention to the risk of adverse cerebrovascular events.<sup>111</sup> They should be avoided or used with extreme caution in patients with DLB, given their sensitivity to these agents. When necessary the second-generation antipsychotics are preferred.<sup>104</sup> If the patient is taking a dopaminergic (anti-parkinsonian) drug, lowering its dose would be the preferred first step before introducing a dopamine-blocking agent.

## Benzodiazepines

In general, benzodiazepines are to be avoided in the neurocognitive disorders because of the risk of paradoxical agitation as well as of falls and further diminished cognition. An exception may be the treatment of REM Sleep Behavior Disorder in DLB.

**Further discussion** of the pharmacologic, psychosocial, and environmental management of neurocognitive disorders is provided elsewhere in this volume.

## SUMMARY

Clinicians should be knowledgeable about the various neurocognitive disorders which are common and devastating in older adults. Diagnosis requires careful history-taking and skilled clinical assessment, followed by appropriate laboratory investigations. Diagnostic imaging can be useful when interpreted by experts familiar with these conditions. Biomarkers for most of these disorders are still being validated and are not yet recommended for clinical use. Referral to specialists can be valuable for specific purposes, e.g., neuropsychologists for objective cognitive testing and interpretation; neurologists for diagnosis, particularly of the less common disorders; geriatric psychiatrists when there are psychological or behavioral challenges. Drug treatments at present provide symptomatic relief. Psychosocial and other supportive therapies are essential.

## References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. Fifth Edition. Arlington, VA: American Psychiatric Association; 2013.
2. Murphy SL, Xu J, Kochanek KD. Division of Vital Statistics. Deaths: final data for 2010. National Vital Statistics Reports. 2013; 61(4) Available at: [http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61\\_04.pdf](http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_04.pdf).
3. Alzheimer's Association. 2013 Alzheimer's disease facts and figures. *Alzheimers Dement.* 2013; 9:208–245. [PubMed: 23507120]
4. Prince M, Bryce R, Albanese E, Wimo A, Ribeiro W, Ferri CP. The global prevalence of dementia: a systematic review and metaanalysis. *Alzheimers Dement.* 2013; 9:63–75. [PubMed: 23305823]
5. Jorm AF, Jolley D. The incidence of dementia: a meta-analysis. *Neurology.* 1998; 51(3):728–733. [PubMed: 9748017]
6. Ward A, Arrighi HM, Michels S, Cedarbaum JM. Mild cognitive impairment: disparity of incidence and prevalence estimates. *Alzheimers Dement.* 2012; 8(1):14–21. [PubMed: 22265588]
7. Prince M, Prina M, Guerchet. Alzheimer's Disease International. World Alzheimer Report 2013 Journey of Caring: An analysis of long term care for dementia. Available at: <http://www.alz.co.uk/research/world-report-2013>.

8. Rocca WA, Petersen RC, Knopman DS, et al. Trends in the incidence and prevalence of Alzheimer's disease, dementia, and cognitive impairment in the United States. *Alzheimers Dement*. 2011; 7(1):80–93. [PubMed: 21255746]
9. Matthews FE, Arthur A, Barnes LE, et al. A two-decade comparison of prevalence of dementia in individuals aged 65 years and older from three geographical areas of England: results of the Cognitive Function and Ageing Study I and II. *Lancet*. 2013; 382(9902):1405–1412. [PubMed: 23871492]
10. Gurland BJ, Wilder DE, Lantigua RF, et al. Rates of dementia in three ethnorracial groups. *International Journal of Geriatric Psychiatry*. 1999; 14(6):481–493. [PubMed: 10398359]
11. Hall KS, Gao S, Baiyewu O, et al. Prevalence rates for dementia and Alzheimer's disease in African Americans: 1992 versus 2001. *Alzheimers Dement*. 2009; 5(3):227–233. [PubMed: 19426950]
12. Plassman BL, Langa KM, Fisher GG, et al. Prevalence of dementia in the United States: the aging, demographics and memory study. *Neuroepidemiology*. 2007; 29:125–132. [PubMed: 17975326]
13. Graves AB, Larson EB, Edland SD, et al. Prevalence of dementia and its subtypes in the Japanese American population of King County, Washington state. *Am J Epidemiol*. 1996; 144(8):760–771. [PubMed: 8857825]
14. Canadian Study of Health and Aging Working Group. Canadian Study of Health and Aging: study methods and prevalence of dementia. *Canadian Medical Association Journal*. 1994; 150(6):899–913. [PubMed: 8131123]
15. Aarsland D, Zaccai J, Brayne C. A symptomatic review of prevalence studies of dementia in Parkinson's disease. *Mov Disord*. 2005; 20(10):1255–1263. [PubMed: 16041803]
16. Zaccai J, McCracken C, Brayne C. A systematic review of prevalence and incidence studies of dementia with Lewy bodies. *Age Ageing*. 2005; 34(6):561–566. [PubMed: 16267179]
17. Rosso SM, Donker Kaat L, Baks T, et al. Frontotemporal dementia in The Netherlands: patient characteristics and prevalence estimates from a population-based study. *Brain*. 2003; 126:2016–2022. [PubMed: 12876142]
18. Gao S, Hendrie HC, Hall KS, Hui S. The relationships between age, sex, and the incidence of dementia and Alzheimer disease: a meta-analysis. *Arch Gen Psychiatry*. 1998; 55(9):809–815. [PubMed: 9736007]
19. Roses AD. Apolipoprotein E alleles as risk factors in Alzheimer's disease. *Annu Rev Med*. 1996; 47:387–400. [PubMed: 8712790]
20. Tsuang D, Leverenz JB, Lopez OL, et al. APO ε4 increases risk for dementia in pure synucleinopathies. *JAMA Neurol*. 2013; 70(2):223–228. [PubMed: 23407718]
21. Chuang Y, Hayden KM, Norton MC, et al. Association between APOE ε4 allele and vascular dementia: the Cache County study. *Dement Geriatr Cogn Disord*. 2010; 29:248–253. [PubMed: 20375505]
22. Yin YW, Li JC, Wang JZ, et al. Association between apolipoprotein E gene polymorphism and the risk of vascular dementia: a meta-analysis. *Neurosci Lett*. 2012; 514(1):6–11. [PubMed: 22381401]
23. Srinivasan R, Davidson Y, Gibbons L, et al. The apolipoprotein E ε4 allele selectively increases the risk of frontotemporal lobar degeneration in males. *J Neurol Neurosurg Psychiatry*. 2006; 77(2):154–158. [PubMed: 16421115]
24. Justin BN, Turek M, Hakim AM. Heart disease as a risk factor for dementia. *Clin Epidemiol*. 2013; 5:135–145. [PubMed: 23658499]
25. Kivipelto M, Ngandu T, Fratiglioni L, et al. Obesity and vascular risk factors at midlife and the risk of dementia and Alzheimer disease. *Arch Neurol*. 2005; 62(10):1556–1560. [PubMed: 16216938]
26. Schnaider Beeri M, Goldbourt U, Silverman JM, et al. Diabetes mellitus in midlife and the risk of dementia three decades later. *Neurology*. 2004; 63(10):1902–1907. [PubMed: 15557509]
27. Tilvis RS, Kähönen-Väre MH, Jolkkonen J, Valvanne J, Pitkala KH, Strandberg TE. Predictors of cognitive decline and mortality of aged people over a 10-year period. *J Gerontol A Biol Sci Med Sci*. 2004; 59(3):268–274. [PubMed: 15031312]

28. Qui C, Winblad B, Marengoni A, Klarin I, Fastbom J, Fratiglioni L. Heart failure and risk of dementia and Alzheimer disease: a population-based cohort study. *Arch Intern Med.* 2006; 166(9): 1003–1008. [PubMed: 16682574]
29. Santangeli P, Di Biase L, Bai R, et al. Atrial fibrillation and the risk of incident dementia: a meta-analysis. *Heart Rhythm.* 2012; 9(11):1761–1768. [PubMed: 22863685]
30. de la Torre JC. Cardiovascular risk factors promote brain hypoperfusion leading to cognitive decline and dementia. *Cardiovasc Psychiatry and Neurol.* 2012 Article ID 367516, 15 pages.
31. Nation DA, Edland SD, Bond MW, et al. Pulse pressure is associated with Alzheimer biomarkers in cognitively normal older adults. *Neurology.* 2013; 81(23):2024–2027. [PubMed: 24225352]
32. Schmidt R, Schmidt H, Curb JD, Masaki K, White LR, Launer LJ. Early inflammation and dementia: a 25-year follow-up of the Honolulu-Asia Aging Study. *Ann Neurol.* 2002; 52(2):168–174. [PubMed: 12210786]
33. Ravaglia G, Forti P, Maioli F, et al. Blood inflammatory markers and risk of dementia: the Conselice Study of Brain Aging. *Neurobiol Aging.* 2007; 28(12):1810–1820. [PubMed: 17011077]
34. Akiyama H, Barger S, Barnum S, et al. Inflammation and Alzheimer's disease. *Neurobiol Aging.* 2000; 21(3):383–421. [PubMed: 10858586]
35. Schott JM, Revesz T. Inflammation in Alzheimer's disease: insights from immunotherapy. *Brain.* 2013; 136(9):2654–2656. [PubMed: 23983027]
36. Tuppo EE, Arias HR. The role of inflammation in Alzheimer's disease. *Int J Biochem Cell Biol.* 2005; 37(2):289–305. [PubMed: 15474976]
37. Budhiraja R, Budhiraja P, Quan SF. Sleep-disordered breathing and cardiovascular disorders. *Respiratory Care.* 2010; 55(10):1322–1332. [PubMed: 20875159]
38. Kim H, Yun CH, Thomas RJ, et al. Obstructive sleep apnea as a risk factor for cerebral white matter change in a middle-aged and older general population. *Sleep.* 2013; 36(5):709–715. [PubMed: 23633753]
39. Chang WP, Liu ME, Chang WC, et al. Sleep apnea and the risk of dementia: a population-based 5-year follow-up study in Taiwan. *PLOS One.* 2013; 8(10):e78655. [PubMed: 24205289]
40. van Kooten F, Koudstaal PJ. Epidemiology of post-stroke dementia. *Haemostasis.* 1998; 28(3–4): 124–133. [PubMed: 10420060]
41. Pendlebury ST, Rothwell PM. Prevalence, incidence, and factors associated with pre-stroke and post-stroke dementia: a systematic review and meta-analysis. *Lancet Neurol.* 2009; 8:1006–1018. [PubMed: 19782001]
42. Dotson VM, Beydoun MA, Zonderman AB. Recurrent depressive symptoms and the incidence of dementia and mild cognitive impairment. *Neurology.* 2010; 75(1):27–34. [PubMed: 20603482]
43. Panza F, Frisardi V, Capurso C, et al. Late-life depression, mild cognitive impairment, and dementia: possible continuum? *Am J Geriatr Psychiatry.* 2010; 18(2):98–116. [PubMed: 20104067]
44. Butters MA, Young JB, Lopez O, et al. Pathways linking late-life depression to persistent cognitive impairment and dementia. *Dialogues Clin Neurosci.* 2008; 10(3):345–357. [PubMed: 18979948]
45. Beaudreau SA, O'Hara R. Late-life anxiety and cognitive impairment: a review. *Am J Geriatr Psychiatry.* 2008; 16(10):790–803. [PubMed: 18827225]
46. Yaffe K, Vittinghoff E, Lindquist K, et al. Posttraumatic stress disorder and risk of dementia among US veterans. *Arch Gen Psychiatry.* 2010; 67(6):608–613. [PubMed: 20530010]
47. Wilson RS, Boyle PA, Buchman AS, Yu L, Arnold SE, Bennett DA. Harm avoidance and risk of Alzheimer's disease. *Psychosom Med.* 2011; 73(8):690–696. [PubMed: 21949425]
48. Fleminger S, Oliver DL, Lovestone S, Rabe-Hesketh S, Giora A. Head injury as a risk factor for Alzheimer's disease: the evidence 10 years on; a partial replication. *J Neurol Neurosurg Psychiatry.* 2003; 74(7):857–862. [PubMed: 12810767]
49. Guo Z, Cupples LA, Kurz A, et al. Head injury and the risk of AD in the MIRAGE study. *Neurology.* 2000; 54:1316–1323. [PubMed: 10746604]

50. Plassman BL, Havlik RJ, Steffens DC, et al. Documented head injury in early adulthood and risk of Alzheimer's disease and other dementias. *Neurology*. 2000; 55:1158–1166. [PubMed: 11071494]
51. Gavett BE, Stern RA, Cantu RD, Nowinski CJ, McKee AC. Mild traumatic brain injury: a risk factor for neurodegeneration. *Alzheimers Res Ther*. 2010; 2(3):18–21. [PubMed: 20587081]
52. Brown RC, Lockwood AH, Sonawane BR. Neurodegenerative diseases: an overview of environmental risk factors. *Environ Health Perspect*. 2005; 113(9):1250–1256. [PubMed: 16140637]
53. Anstey KJ, von Sanden C, Salim A, O'Kearney R. Smoking as a risk factor for dementia and cognitive decline: a meta-analysis of prospective studies. *Am J Epidemiol*. 2007; 166(4):367–378. [PubMed: 17573335]
54. Chang CC, Zhao Y, Lee CW, Ganguli M. Smoking, death, and Alzheimer disease: a case of competing risks. *Alzheimer Dis Assoc Disord*. 2012; 26(4):300–306. [PubMed: 22185783]
55. Quik M, Perez XA, Bordia T. Nicotine as a potential neuroprotective agent for Parkinson's disease. *Mov Disord*. 2012; 27(8):947–957. [PubMed: 22693036]
56. Anttila T, Helkala EL, Viitanen M, et al. Alcohol drinking in middle age and subsequent risk of mild cognitive impairment and dementia in old age: a prospective population based study. *BMJ*. 2004; 329(7465):539. [PubMed: 15304383]
57. Mukamal KJ, Kuller LH, Fitzpatrick AL, Longstreth WT Jr, Mittleman MA, Siscovick DS. Prospective study of alcohol consumption and risk of dementia in older adults. *JAMA*. 2003; 289(11):1405–1413. [PubMed: 12636463]
58. Franco R, Li S, Rodriguez-Rocha H, Burns M, Panayiotidis MI. Molecular mechanisms of pesticide-induced neurotoxicity: Relevance to Parkinson's disease. *Chem Biol Interact*. 2010; 188(2):289–300. [PubMed: 20542017]
59. Fratiglioni L, Wang HX. Brain reserve hypothesis in dementia. *J Alzheimers Dis*. 2007; 12(1):11–22. [PubMed: 17851191]
60. Katzman R, Terry R, DeTeresa R, et al. Clinical, pathological, and neurochemical changes in dementia: a subgroup with preserved mental status and numerous neocortical plaques. *Ann Neurol*. 1988; 23(2):138–144. [PubMed: 2897823]
61. Stern Y. What is cognitive reserve? Theory and research application of the reserve concept. *J Int Neuropsychol Soc*. 2002; 8(3):448–460. [PubMed: 11939702]
62. Meng X, D'Arcy C. Education and dementia in the context of the cognitive reserve hypothesis: a systematic review with meta-analyses and qualitative analyses. *PLoS One*. 2012; 7(6):e38268. [PubMed: 22675535]
63. Alladi S, Bak T, Duggirala V, et al. Bilingualism delays age at onset of dementia, independent of education and immigrations status. *Neurology*. 2013; 81(2):1938–1944. [PubMed: 24198291]
64. Craik FI, Bialystok E, Freedman M. Delaying the onset of Alzheimer disease: bilingualism as a form a cognitive reserve. *Neurology*. 2010; 75(19):1726–1729. [PubMed: 21060095]
65. Bonaiuto S, Rocca WA, Lippi A, et al. Education and occupation as risk factors for dementia: a population-based case control study. *Neuroepidemiology*. 1995; 14(3):101–109. [PubMed: 7777124]
66. Bickel H, Kurz A. Education, occupation, and dementia: the Bavarian school sisters study. *Dement Geriatr Cogn Disord*. 2009; 27(6):548–556. [PubMed: 19590201]
67. Verghese J, Lipton RB, Katz MJ, et al. Leisure activities and the risk of dementia in the elderly. *N Engl J Med*. 2003; 348(25):2508–2516. [PubMed: 12815136]
68. Hughes TF. Promotion of cognitive health through activity in the aging population. *Aging Health*. 2010; 6(1):111–121. [PubMed: 20383318]
69. Yip AG, Green RC, Huyck M, Cupples LA, Farrer LA. MIRAGE Study Group. Nonsteroidal anti-inflammatory drug use and Alzheimer's disease risk: the MIRAGE study. *BMC Geriatrics*. 2005; 5:2. [PubMed: 15647106]
70. Szekely CA, Breitner JC, Fitzpatrick AL, et al. NSAID use and dementia risk in the Cardiovascular Health Study: role of APOE and NSAID type. *Neurology*. 2008; 70(1):17–24. [PubMed: 18003940]

71. in't Veld BA, Launer LJ, Hoes AW, et al. NSAIDs and incident Alzheimer's disease. The Rotterdam Study. *Neurobiol Aging*. 1998; 19(6):607–611. [PubMed: 10192221]
72. de Craen AJ, Gussekloo J, Vrijnsen B, Westendorp RG. Meta-analysis of nonsteroidal anti-inflammatory drug use and risk of dementia. *Am J Epidemiol*. 2005; 161(2):114–120. [PubMed: 15632261]
73. Swiger KJ, Manalac RJ, Blumenthal RS, Blaha MJ, Martin SS. Statins and cognition: a systematic review and meta-analysis of short- and long-term cognitive effects. *Mayo Clin Proc*. 2013; 88(11): 1213–1221. [PubMed: 24095248]
74. Wong WB, Lin VW, Boudreau D, Devine EB. Statins in the prevention of dementia and Alzheimer's disease: a meta-analysis of observational studies and an assessment of confounding. *Pharmacoeconomics Drug Saf*. 2013; 22(4):345–358.
75. Shumaker SA, Legault C, Rapp SR, et al. Estrogen plus progestin and the incidence of dementia and mild cognitive impairment in postmenopausal women: the Women's Health Initiative Memory Study: a randomized controlled trial. *JAMA*. 2003; 289(20):2651–2662. [PubMed: 12771112]
76. Shumaker SA, Legault C, Kuller L, et al. Conjugated equine estrogens and incidence of probable dementia and mild cognitive impairment in postmenopausal women: Women's Health Initiative Memory Study. *JAMA*. 2004; 291(24):2947–2958. [PubMed: 15213206]
77. O'Brien J, Jackson JW, Grodstein F, Blacker D, Weuve J. Postmenopausal hormone therapy is not associated with risk of all-cause dementia and Alzheimer's Disease. *Epidemiol Rev*. first published online September 15, 2013.
78. Shao H, Breitner JC, Whitmer RA, et al. Hormone therapy and Alzheimer disease dementia: new findings from the Cache County Study. *Neurology*. 2012; 79(18):1846–1852. [PubMed: 23100399]
79. Whitmer RA, Quesenberry CP, Zhou J, Yaffe K. Timing of hormone therapy and dementia: the critical window theory revisited. *Ann Neurol*. 2011; 69(1):163–169. [PubMed: 21280086]
80. Zandi PP, Carlson MC, Plassman BL, et al. Hormone replacement therapy and incidence of Alzheimer disease in older women: the Cache County Study. *JAMA*. 2002; 288(17):2123–2129. [PubMed: 12413371]
81. Ruitenberg A, van Swieten JC, Witteman JC, et al. Alcohol consumption and risk of dementia: the Rotterdam study. *Lancet*. 2002; 359(9303):281–286. [PubMed: 11830193]
82. Ganguli M, Vander Bilt J, Saxton JA, Shen C, Dodge HH. Alcohol consumption and cognitive function in late life: a longitudinal community study. *Neurology*. 2005; 65(8):1210–1217. [PubMed: 16247047]
83. Lourida I, Soni M, Thompson-Coon J, et al. Mediterranean diet, cognitive function, and dementia: a systematic review. *Epidemiology*. 2013; 24(4):479–489. [PubMed: 23680940]
84. Hamer M, Chida Y. Physical activity and risk of neurodegenerative disease: a systematic review of prospective evidence. *Psychol Med*. 2009; 39(1):3–11. [PubMed: 18570697]
85. Crooks VC, Lubben J, Petitti DB, Little D, Chiu V. Social network, cognitive function, and dementia incidence among elderly women. *Am J Public Health*. 2008; 98(7):1221–1227. [PubMed: 18511731]
86. Wang HX, Karp A, Winblad B, Fratiglioni L. Late-life engagement in social and leisure activities is associated with a decreased risk of dementia: a longitudinal study from the Kungsholmen Project. *Am J Epidemiol*. 2002; 155(12):1081–1087. [PubMed: 12048221]
87. Fratiglioni L, Paillard-Bog S, Winblad B. An active and socially integrated lifestyle in late life might protect against dementia. *Lancet Neurol*. 2004; 3(6):343–353. [PubMed: 15157849]
88. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state." A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12(3):189–198. [PubMed: 1202204]
89. Nasreddine ZS, Phillips NA, Bédirian V. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc*. 2005; 53(4):695–699. [PubMed: 15817019]
90. Borson S, Scanlan J, Brush M, Vitaliano P, Dokmak A. The mini-cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry*. 2000; 15(11): 1021–1027. [PubMed: 11113982]

91. Brookmeyer R, Corrada MM, Curriero FC, Kawas C. Survival following a diagnosis of Alzheimer disease. *Arch Neurol*. 2002; 59(11):1764–1767. [PubMed: 12433264]
92. Helzner EP, Scarmeas N, Cosentino S, Tang MX, Schupf N, Stern Y. Survival in Alzheimer disease: a multiethnic, population-based study of incident cases. *Neurology*. 2008; 71(19):1489–1495. [PubMed: 18981370]
93. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging- Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011; 7(3): 263–269. [PubMed: 21514250]
94. Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging- Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011; 7(3): 280–292. [PubMed: 21514248]
95. Jellinger KA. Morphological diagnosis of “vascular dementia”- a critical update. *J Neurol Sci*. 2008; 270(1–2):1–12. [PubMed: 18455191]
96. Sneed JR, Culang-Reinlieb ME. The vascular depression hypothesis: an update. *Am J Geriatr Psychiatry*. 2011; 19(2):99–103. [PubMed: 21328801]
97. Rabinovici GD, Miller BL. Frontotemporal lobar degeneration: epidemiology, pathophysiology, diagnosis and management. *CNS Drugs*. 2010; 24(5):375–398. [PubMed: 20369906]
98. McKeith IG, Dickson DW, Lowe J, et al. Diagnosis and management of dementia with Lewy bodies: third report of the DLB consortium. *Neurology*. 2005; 65(12):1863–1872. [PubMed: 16237129]
99. Brown K, Mastrianni JA. The prion diseases. *J Geriatr Psychiatry Neurol*. 2010; 23(4):277–298. [PubMed: 20938044]
100. Birks J. Cholinesterase inhibitors for Alzheimer's disease. *Cochrane Database of Systematic Reviews*. 2006; (1):CD005593. [PubMed: 16437532]
101. Press, D.; Alexander, M. Cholinesterase inhibitors in the treatment of dementia. In: Basow, DS., editor. *UpToDate*. Waltham, MA: UpToDate; [Accessed on December 17, 2013]
102. Emre M, Aarsland D, Albanese A, et al. Rivastigmine for dementia associated with Parkinson's disease. *N Engl J Med*. 2004; 351(24):2509–2518. [PubMed: 15590953]
103. Maidment I, Fox C, Boustani M. Cholinesterase inhibitors for Parkinson's disease dementia. *Cochrane Database of Systematic Reviews*. 2006; (1):CD004747. [PubMed: 16437494]
104. McKeith I, Mintzer J, Aarsland D, et al. Dementia with Lewy Bodies. *Lancet Neurology*. 2004; 3(1):19–28. [PubMed: 14693108]
105. Wright, CB. Treatment and prevention of vascular dementia. In: Basow, DS., editor. *UpToDate*. Waltham, MA: UpToDate; [Accessed on December 17, 2013]
106. Miller, BL.; Lee, SE. Frontotemporal dementia: treatment. In: Basow, DS., editor. *UpToDate*. Waltham, MA: UpToDate; [Accessed on December 17, 2013]
107. Mendez MF, Shapira JS, McMurtray A, Licht E. Preliminary findings: behavioral worsening on donepezil in patients with frontotemporal dementia. *Am J Geriatr Psychiatry*. 2007; 15(1):84–87. [PubMed: 17194818]
108. Russ TC, Morling JR. Cholinesterase inhibitors for mild cognitive impairment. *Cochrane Database of Systematic Reviews*. 2012; 9:CD009132. [PubMed: 22972133]
109. McShane R, Areosa Sastre A, Minakaran N. Memantine for dementia. *Cochrane Database of Systematic Reviews*. 2006; 2:CD003154. [PubMed: 16625572]
110. Huey ED, Putnam KT, Grafman J. A systematic review of neurotransmitter deficits and treatments in frontotemporal dementia. *Neurology*. 2006; 66(1):17–22. [PubMed: 16401839]
111. Gareri P, DeFazio P, Manfredi VG, DeSarro G. Use and safety of antipsychotics in behavioral disorders in elderly people with dementia. *J Clin Psychopharmacol*. Published Ahead-of-Print.

### Key Points

1. Clinicians should be knowledgeable about the various neurocognitive disorders which are common and devastating in older adults.
2. Diagnosis requires careful history-taking and skilled clinical assessment, followed by appropriate laboratory investigations.
3. Diagnostic imaging can be useful when interpreted by experts familiar with these conditions.
4. Biomarkers for most of these disorders are still being validated and are not yet recommended for clinical use.
5. Referral to specialists can be valuable for specific purposes, e.g., neuropsychologists for objective cognitive testing and interpretation; neurologists for diagnosis, particularly of the less common disorders; geriatric psychiatrists when there are psychological or behavioral challenges.
6. Drug treatments at present provide symptomatic relief. Psychosocial and other supportive therapies are essential.

**Table 1**

Neurocognitive Disorders as Diagnosed in DSM-5

Diagnostic Criteria	Major Neurocognitive Disorder/Dementia	Major Neurocognitive Disorder/Dementia
A	Significant cognitive decline in one or more cognitive domains, based on:	Modest cognitive decline in one or more cognitive domains, based on:
	1. Concern about significant decline, expressed by individual or reliable informant, or observed by clinician.	1. Concern about mild decline, expressed by individual or reliable informant, or observed by clinician.
	2. Substantial impairment, documented by objective cognitive assessment.	2. Modest impairment, documented by objective cognitive assessment.
B	Interference with independence in everyday activities.	No interference with independence in everyday activities, although these activities may require more time and effort, accommodation, or compensatory strategies
C	Not exclusively during delirium.	
D	Not better explained by another mental disorder.	
E	Specify one or more etiologic subtypes, "due to"	
	<ul style="list-style-type: none"> <li>• Alzheimer's disease</li> <li>• Cerebrovascular disease (Vascular Neurocognitive Disorder)</li> <li>• Frontotemporal Lobar Degeneration (Frontotemporal Neurocognitive Disorder)</li> <li>• Dementia with Lewy Bodies (Neurocognitive Disorder with Lewy Bodies)</li> <li>• Parkinson's disease</li> <li>• Huntington's disease</li> <li>• Traumatic Brain Injury</li> <li>• HIV Infection</li> <li>• Prion Disease</li> <li>• Another medical condition</li> <li>• Multiple etiologies</li> </ul>	

*Adapted from:* American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association; 2013

Table 2

Prevalence of Dementia: Overall and Subtypes

Authors	Country	Study type	Age	Regions	Overall Dementia	Alzheimer'S dementia	Vascular dementia	Parkinson'S dementia	Dementia with Lewy Bodies	Fronto-temporal dementia	
Prince <i>et al.</i> , 2013 <sup>4</sup>	Global	Meta-analysis + systematic review	>=60	Latin America	8.5%						
				Sub-Saharan Africa	5.0–7.0%						
				Other world regions	2.0–4.0%						
*Matthews <i>et al.</i> , 2013 <sup>9</sup>	UK	Population survey	65+	Cambridgeshire, Newcastle, and Nottingham	6.5%						
				Hispanic/Latino	65–74	7.5%					
					75–84	27.9%					
Gurland <i>et al.</i> , 1999 <sup>10</sup>	USA	Population survey	65+	African American	85+	62.9%					
					65–74	9.1%					
					75–84	19.9%					
				Non-Hispanic White	85+	58.6%					
					65–74	2.9%					
					75–84	10.9%					
					85+	30.2%					
*Hall <i>et al.</i> , 2009 <sup>11</sup>	USA	Population survey	70+	African Americans in Indianapolis	7.45%	6.77%					
Plassman <i>et al.</i> , 2007 <sup>12</sup>	USA	Population survey	71–79	Nationally representative sample	4.97%	2.32%	0.98%				
			80–89		24.19%	18.1%	4.09%				
			90+		37.36%	29.7%	6.19%				
*Graves <i>et al.</i> , 1996 <sup>13</sup>	USA	Population survey	65+	Japanese Americans	6.32%	3.46%	1.41%				
*CSHA Working Group, 1994 <sup>14</sup>	Canada	Population survey	65+	Nationally representative sample	8%	5.1%	1.5%				
Aarsland <i>et al.</i> , 2005 <sup>15</sup>	Multi-national	Systematic review	65+				0.15–0.5%				
Zaccai <i>et al.</i> , 2005 <sup>16</sup>	Multi-national	Systematic review	65+					0–5%			
Rosso <i>et al.</i> , 2003 <sup>17</sup>	Netherlands	Population survey	50–59						Per 100,000		
										3.6	

Authors	Country	Study type	Age		Regions	Overall Dementia	Alzheimer's dementia	Vascular dementia	Parkinson's dementia	Dementia with Lewy Bodies	Fronto-temporal dementia	
			60-69	70-79								
											9.4	
											3.8	

In all studies, prevalence % increases with age.

\* Age-specific prevalence is reported in the original articles but omitted from this table.

**Table 3**

Functional limitations associated with impairment in different cognitive domains

<b>Cognitive domain</b>	<b>Examples of changes in everyday activities</b>
Complex attention	Normal tasks take longer, especially when there are competing stimuli; easily distracted; tasks need to be simplified; difficulty holding information in mind to do mental calculations or dial a phone number
Executive functioning	Difficulty with multi-stage tasks, planning, organizing, multi-tasking, following directions, keeping up with shifting conversations
Learning and memory	Difficulty recalling recent events, repeating self, misplacing objects, losing track of actions already performed, increasing reliance on lists, reminders
Language	Word-finding difficulty, use of general phrases or wrong words, grammatical errors, difficulty with comprehension of others' language or written material
Perceptual-motor/visuospatial function	Getting lost in familiar places, more use of notes and maps, difficulty using familiar tools and appliances
Social cognition	Disinhibition or apathy, loss of empathy, inappropriate behavior, loss of judgment

*Adapted from:* American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association; 2013

**Table 4**

Examples of objective cognitive assessments, as noted in DSM-5

Cognitive domains	Objective Assessment
Complex Attention	<i>Maintenance of attention</i> , e.g., press a button every time a tone is heard, over a period of time. <i>Selective attention</i> , e.g., hear numbers and letters, but count only the letters. <i>Divided attention</i> , e.g., tap rapidly while learning a story. <i>Processing speed</i> : carry out any timed task.
Executive Functioning	<i>Planning</i> : e.g., maze puzzles, interpret sequential pictures or arrange objects in sequence. <i>Decision making</i> with competing alternatives, e.g., simulated gambling. <i>Working memory</i> : hold information for a brief period and manipulate it, e.g. repeat a list of numbers backward. <i>Feedback utilization</i> : Use feedback on errors to infer rules to carry out tasks. <i>Inhibition</i> : Override habits; choose the correct but more complex and less obvious solution, e.g., read printed names of colors rather than naming the color in which they are printed. <i>Cognitive flexibility</i> : Shift between sets, concepts, tasks, rules, e.g., alternate between numbers and letters.
Learning and Memory	<i>Immediate memory</i> : Repeat a list of words or digits. <i>Recent memory</i> : Free recall: recall as many items as possible from, e.g., a list of words, or a story, or a diagram. Cued recall: with examiner providing cues, e.g., “recall as many food items as you can from the list.” Recognition: with examiner asking, e.g., “was there an apple on the list?” <i>Semantic memory</i> : recall well-known facts. <i>Autobiographical memory</i> : recall personal events. <i>Implicit (procedural) memory</i> : recall skills to carry out procedures.
Language	<i>Expressive language</i> : confrontation naming of e.g., objects or pictures; fluency for words in a given category (e.g. animals) or beginning with a given letter, as many as possible in one minute. <i>Grammar and syntax</i> : omitting or incorrectly using articles, prepositions, helper verbs. <i>Receptive language</i> : comprehend /define words, carry out simple commands.
Perceptuomotor functioning	<i>Visuoconstructional</i> : e.g., Draw, copy, assemble blocks. <i>Perceptuomotor</i> : e.g., Insert blocks or pegs into appropriate slots. <i>Praxis</i> : Mime gestures such as “salute” or actions such as “use hammer.” <i>Gnosis</i> : e.g., recognize faces and colors.
Social cognition	<i>Recognize emotions</i> : Identify pictures showing e.g., happy, sad, scared, angry faces. <i>Theory of mind</i> : Consider another person’s thoughts, intentions when looking at story cards, e.g., “why is the boy sad?”

Adapted from: American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association; 2013